

Circulatory Mechanisms Underlying Adaptive Increases in Thermogenic Capacity in High-Altitude Deer Mice

Kevin B. Tate^{1,3,4}, Catherine M. Ivy¹, Jonathan P. Velotta², Jay F. Storz³, Grant B. McClelland¹, Zachary A. Cheviron², and Graham R. Scott^{1,*}

¹ Department of Biology, McMaster University, Hamilton, ON, Canada

² Division of Biological Sciences, University of Montana, Missoula, MT, USA

³ School of Biological Sciences, University of Nebraska, Lincoln, NE, USA

⁴ Current address: Department of Biology, Truman State University, Kirksville, MO, USA

* Author for correspondence (scottg2@mcmaster.ca)

Key words: Evolutionary physiology, high-altitude adaptation, respiration, O₂ transport pathway, aerobic performance

Abstract

We examined the circulatory mechanisms underlying adaptive increases in thermogenic capacity in deer mice (*Peromyscus maniculatus*) native to the cold hypoxic environment at high altitudes. Deer mice from high- and low-altitude populations were born and raised in captivity to adulthood, and then acclimated to normoxia or hypobaric hypoxia (simulating hypoxia at ~4300 m). Thermogenic capacity (maximal O₂ consumption, VO₂max, during cold exposure) was measured in hypoxia, along with arterial O₂ saturation (SaO₂) and heart rate (f_H). Hypoxia acclimation increased VO₂max by a greater magnitude in highlanders than in lowlanders. Highlanders also had higher SaO₂ and extracted more O₂ from the blood per heartbeat (O₂ pulse = VO₂max/f_H). Hypoxia acclimation increased f_H, O₂ pulse, and capillary density in the left ventricle of the heart. Our results suggest that adaptive increases in thermogenic capacity involve integrated functional changes across the O₂ cascade that augment O₂ circulation and extraction from the blood.

Summary Statement

Adaptive increases in thermogenic capacity in high-altitude deer mice involve integrated functional changes across the O₂ cascade that augment O₂ circulation and extraction from the blood.

INTRODUCTION

High-altitude natives are valuable model organisms to understand how physiological systems evolve. The cold and oxygen-depleted ('hypoxic') environment at high altitudes requires that endotherms sustain high rates of O₂ consumption for thermogenesis and locomotion while facing a diminished O₂ supply. Growing evidence suggests that high-altitude natives have overcome this challenge through evolved changes in the physiological systems underlying O₂ transport and utilization (Monge and León-Velarde, 1991; Storz et al., 2010b; Scott, 2011; Ivy and Scott, 2015). Studies of high-altitude natives aimed at understanding the evolution of the O₂ transport cascade – comprised of ventilation, pulmonary diffusion, circulation, tissue diffusion, and cellular O₂ utilization – are therefore extremely valuable for explaining the physiological mechanisms of evolutionary adaptation.

North American deer mice (*Peromyscus maniculatus*) are an excellent model species for studies of high-altitude adaptation. Their native altitudinal range extends from below sea level in Death Valley California to over 4300 m above sea level in the Rocky Mountains (Hock, 1964; Snyder et al., 1982; Natarajan et al., 2015). High-altitude populations must sustain high metabolic rates in the wild (Hayes, 1989b), and there appears to be strong directional selection on thermogenic capacity (maximal O₂ consumption, VO₂max, during cold exposure) to support heat generation in cold alpine environments (Hayes and O'Connor, 1999). In response to this strong selection pressure, high-altitude deer mice have evolved a higher VO₂max in hypoxia than low-altitude populations of deer mice and a congeneric lowland species (white-footed mice, *P. leucopus*) (Cheviron et al., 2012; Cheviron et al., 2013; Lui et al., 2015). This evidence suggests that highland deer mice have evolved an adaptive increase in thermogenic capacity in hypoxia.

The physiological mechanisms underlying this evolved increase in thermogenic capacity have yet to be fully explained. High-altitude deer mice have evolved a high blood-O₂ affinity compared to their lowland counterparts that contributes to increasing VO₂max in hypoxia (Snyder et al., 1982; Chappell and Snyder, 1984; Storz et al., 2010a; Natarajan et al., 2013), but it is unclear if this adaptation improves O₂ uptake into the blood *in vivo*. High-altitude deer mice have also evolved a more oxidative and richly vascularized phenotype of the skeletal muscle (used for shivering and locomotion), in association with differential expression of genes involved in aerobic energy metabolism and angiogenesis (Cheviron et al., 2012; Cheviron et al., 2014; Lui et al., 2015; Scott et al., 2015; Lau et al., 2017; Mahalingam et al., 2017). Development and

acclimatization to cold/hypoxia are also known to affect VO_2max , cardiopulmonary organ sizes, and the capacity for non-shivering thermogenesis in deer mice (Hammond et al., 2001; Hammond et al., 2002; Chappell and Hammond, 2004; Shirkey and Hammond, 2014; Velotta et al., 2016). However, we know very little about *in vivo* cardiorespiratory function at VO_2max in this species. This study therefore aims to examine the contribution of differences in arterial O_2 saturation and some other aspects of circulatory function to adaptive increases in thermogenic capacity in high-altitude deer mice.

MATERIALS AND METHODS

Animals and acclimation treatments

Captive breeding populations were established from wild deer mouse populations native to high altitude (near the summit of Mount Evans, CO, USA at 39°35'18"N, 105°38'38"W; 4,350 m above sea level) (*P. m. rufinus*) and low altitude (Nine Mile Prairie, Lancaster County, NE, USA at 40°52'12"N, 96°48'20.3"W; 430 m above sea level) (*P. m. nebracensis*). Wild adults were transported to McMaster University (near sea level) and housed in common-garden conditions, and were bred within each population to produce lab-raised progeny. Mice were raised in standard holding conditions (24-25°C, 12 h light: 12 h dark photoperiod) with unlimited access to standard rodent chow and water. All animal protocols followed guidelines established by the Canadian Council on Animal Care and were approved by the McMaster University Animal Research Ethics Board.

Adult mice were raised to ~6 months of age, and a randomly selected group of individuals (mix of both sexes) from each population were acclimated to either (i) normobaria in standard normoxic conditions, or (ii) hypobaric hypoxia (barometric pressure of 60 kPa; equivalent to that at an elevation of ~4300 m) in specially designed hypobaric chambers that have been described previously (Lui et al., 2015). Cages were cleaned twice a week during acclimations, which required that the hypobaric groups be returned to normobaria for a brief period (<30 min). Mice were subjected to subsequent measurements after 6-8 weeks of acclimation.

Respirometry and pulse oximetry

We measured thermogenic capacity in hypoxia in second-generation (F_2) mice from high-altitude and low-altitude populations. Maximal rates of O_2 consumption (VO_{2max}) were measured during acute cold exposure, using open-flow respirometry in a hypoxic heliox atmosphere (12% O_2 , 88% He) at $-5^\circ C$ (Rosenmann and Morrison, 1974; Chappell and Hammond, 2004; Cheviron et al., 2012). Respirometry was carried out in a 0.5 l animal chamber that received a constant incurrent flow rate of 1000 ml/min, regulated using a mass flow controller (MFC-4, Sable Systems) and a precision flow control valve that was factory calibrated for heliox (Sierra Instruments). The chamber was held inside a freezer, in which the ambient temperature was regulated at or slightly below $-5^\circ C$ (measured with a thermocouple; PT-6, Physitemp), and the incurrent gas flowed through copper coils before entering the chamber. Excurrent gas was subsampled at 200 ml/min, dried with pre-baked Drierite, and analyzed for O_2 and CO_2 fractions (FoxBox Respirometry System, Sable Systems).

Respirometry experiments were carried out as follows. Baseline O_2 and CO_2 fractions were first measured without an animal in the chamber. Mice were instrumented with a collar sensor to measure heart rate (f_H) and the O_2 saturation of arterial blood (SaO_2) using a MouseOx Plus pulse oximeter (Starr Life Sciences), and were then transferred to the chamber. The pulse oximetry measurements required that hair be removed from around the neck, which was done two days before the experiments using NairTM hair removal product. Incurrent gas flow rate, chamber temperature, and excurrent O_2 and CO_2 fractions were measured continuously and were acquired using a PowerLab 8/32 and Labchart 8 Pro software (ADInstruments). Pulse oximetry measurements were recorded using Starr Life Sciences acquisition software. Rates of O_2 consumption (VO_2) were calculated using established formulas (Lighton, 2008) and VO_{2max} was defined as the highest VO_2 achieved over a 30s period during the trial, which generally occurred after ~4-6 min in the chamber, when maximal values of SaO_2 and f_H were also determined. Measurements of core body temperature were made using a rectal probe (RET-3-ISO, Physitemp) immediately after removing the animal from the chamber (after ~10-12 min in the chamber), and confirmed that all mice were hypothermic at the end of the experiment.

Cardiac histology

Capillarity was measured histologically in the left ventricle of the heart in a separate group of F₁ mice from highland and lowland populations. Mice were euthanized with an overdose of isoflurane followed by cervical dislocation. The ventricles were removed, coated in embedding medium, frozen in liquid N₂-cooled isopentane, and stored at -80°C. Tissue was sectioned (10 μm) perpendicular to the long axis of the heart in a cryostat at -20°C. Capillaries were identified by staining for alkaline phosphatase activity for 1 h at room temperature (assay buffer concentrations in mM: 1.0 nitroblue tetrazolium, 0.5 5-bromo-4-chloro-3-indoxyl phosphate, 28 NaBO₂, and 7 MgSO₄; pH 9.3). Images were collected systematically using light microscopy, such that there was equal representation of images from across the left ventricle. A blind observer determined the average value of capillary density for each individual.

Statistics

Two-factor ANOVA was generally used to assess the main effects of population altitude and acclimation environment (interactions were also assessed, but were not generally significant and are not reported). Data for VO₂max and the amount of O₂ extracted from the blood per heartbeat (the quotient of VO₂ and f_H; also called the O₂ pulse) were first corrected for body mass (M_b) before making statistical comparisons. This was accomplished by carrying out least-squares regressions to the equation $Y = a M_b^b$ (using GraphPad Prism software; La Jolla, CA), including all of the data across all groups, and then calculating the residual from the regression for each individual. These residuals were then used in two-factor ANOVA, and are reported graphically on the right y-axis. The scale of the left y-axis for graphs of our VO₂max and O₂ pulse data shows the sum of the residual and the expected value for an average-sized 21.6-g mouse (i.e., VO₂max or O₂ pulse data corrected to a body mass of 21.6 g). We also performed a supplementary statistical analysis of the effects of body mass, population altitude, and acclimation environment on VO₂max and O₂ pulse using linear models (lm) in R (R Core Team, 2016), in which body mass and the variable of interest were log-transformed before making statistical comparisons (the statistical results are extremely similar to those obtained with two-factor ANOVA, and are shown in Table S1). Data are generally reported as mean ± s.e.m.

(except when data points from individual samples are shown). $P < 0.05$ was considered significant.

RESULTS AND DISCUSSION

Thermogenic capacity in hypoxia is enhanced in high-altitude deer mice

Cold-induced VO_2max (measured in hypoxia) was greatest in high-altitude deer mice, as reflected by a significant population effect in two-factor ANOVA (Fig. 1A). This was particularly apparent after hypoxia acclimation, when VO_2max was 16% higher on average in highlanders than in lowlanders. Because we observed allometric rather than isometric scaling of VO_2max to body mass, we used a residual-based approach to correct for body mass before making these comparisons (Fig. 1B), but we obtained very similar statistical results using linear model statistics (Table S1). Body mass was similar between populations ($F_{1,36}=1.08$, $p=0.31$) and between acclimation environments ($F_{1,36}=0.49$, $p=0.49$) (normoxic highlanders, 22.9 ± 1.1 g; hypoxic highlanders, 21.0 ± 0.6 g; normoxic lowlanders, 20.6 ± 0.8 g; hypoxic lowlanders, 20.9 ± 1.3 g).

Our results are consistent with previous findings in deer mice and other high-altitude taxa. The increases in cold- and exercise-induced VO_2max in highland deer mice observed by us and others appear to be greatest in hypoxic conditions, and are not as large in normoxic conditions at sea level, suggesting that highlanders are more resistant to the depressing effects of hypoxia on O_2 transport (Chappell and Snyder, 1984; Hayes, 1989a; Cheviron et al., 2012; Cheviron et al., 2013; Lui et al., 2015). Similar differences exist in Andean and Tibetan human populations, in which exercise-induced VO_2max is only elevated compared to lowland humans when tested at altitudes above ~ 2500 m (Brutsaert, 2016). However, in many of these human studies, it has been hard to distinguish evolved genetic effects from effects of developmental environment and exercise training. Although hypoxia exposure during development also has a strong influence on VO_2max in deer mice, directional selection on VO_2max at high altitudes appears to have further increased VO_2max in high-altitude populations (Fig. 1) (Hayes and O'Connor, 1999; Chappell et al., 2007; Russell et al., 2008; Cheviron et al., 2013; Lui et al., 2015).

High-altitude deer mice maintain higher arterial O₂ saturation in hypoxia

Arterial O₂ saturation was ~6-8% higher in highlanders than in lowlanders at VO₂max in hypoxia (Fig. 2A). This observation likely results at least in part from the greater blood- and haemoglobin-O₂ affinities of highlanders (Snyder et al., 1982; Storz et al., 2010a), which would increase SaO₂ at similar conditions of blood O₂ and CO₂ tensions and pH. This observation could also stem from population differences in arterial O₂ tension, arising from differences in pulmonary ventilation or O₂ diffusion. Breathing and pulmonary O₂ extraction have yet to be examined in deer mice at VO₂max, but there appear to be evolved differences in control of breathing by hypoxia under routine conditions in highland deer mice (Ivy and Scott, 2017).

SaO₂ was unaffected by hypoxia acclimation (Fig. 2A), and was not always associated with clear population differences in VO₂max (Fig. 1). Previous studies using wild-derived strains of deer mice with distinct α -globin haplotypes (on randomized genetic backgrounds) have shown that variation in blood-O₂ affinity affects VO₂max, such that mice with higher affinity (typical of highland populations) had the highest VO₂max when acclimated and tested at high altitude (Chappell and Snyder, 1984; Chappell et al., 1988). This relationship is presumed to arise from a positive association between blood-O₂ affinity and SaO₂ in hypoxia, but this has not been tested. Here, the higher SaO₂ in highlanders compared to lowlanders only appears to be associated with increases in VO₂max when mice were acclimated and tested in hypoxia (Fig. 1). However, in normoxia acclimated mice, highlanders had higher SaO₂ without any clear difference in hypoxic VO₂max. This suggests that the influence of SaO₂ on VO₂max may be context dependent, such that the relative benefit of increases in SaO₂ may depend upon interactions with other respiratory traits that change after hypoxia acclimation.

Differences in cardiac performance appear to underlie differences in thermogenic capacity

Heart rates (f_H) during VO₂max in hypoxia were ~9-23% higher after hypoxia acclimation (Fig. 2B). The amount of O₂ extracted from the blood per heartbeat ('O₂ pulse', quotient of VO₂ and f_H) increased by ~25-32% after hypoxia acclimation, and was 10-16% greater in the highland population (Fig. 2C, Table S1). The latter observation suggests that cardiac stroke volume (V_S) and/or the absolute O₂ extraction from the blood ($CaO_2 - CvO_2$) contributes to the variation in VO₂max. This is because all of the above variables are related by the Fick equation, $VO_2 = f_H \times$

$V_S \times (CaO_2 - CvO_2)$, such that O_2 pulse is equal to the product of stroke volume and blood O_2 extraction. This product must therefore be greater in highlanders and increase with hypoxia acclimation.

The observed difference in cardiac performance was likely associated with variation in O_2 supply to heart tissue. Hypoxia acclimation increased capillary density – a key determinant of O_2 diffusing capacity – by ~10-12% in the left ventricle (Fig. 3). However, capillary densities were similar between highlanders and lowlanders, so this trait does not underlie population differences in cardiac performance. Nevertheless, it is likely that an interaction between the hypoxia-induced increase in heart capillarity and the population difference in SaO_2 resulted in an improved O_2 supply to cardiac tissue, and may therefore account for the observed differences in cardiac performance and VO_{2max} . High-altitude adaptation and/or hypoxia acclimation could have also improved the heart's ability to maintain cardiac output during tissue hypoxia. In support of this possibility, some other high-altitude taxa exhibit differences in mitochondrial physiology and metabolic capacity that could improve cardiac function at low O_2 tensions (Sheafor, 2003; Scott et al., 2011; Dawson et al., 2016).

The functional mechanisms of high-altitude adaptation span the O_2 cascade

A key goal of evolutionary physiology is to elucidate the mechanistic basis of adaptive variation in organismal performance (Garland and Carter, 1994; Dalziel et al., 2009). Thermogenesis is a key performance trait that is critical for fitness in endotherms at high altitudes (Hayes and O'Connor, 1999) and can push the respiratory system of many small mammals to its limits (Rosenmann and Morrison, 1974; Chappell and Hammond, 2004). Here, we contribute to the growing evidence suggesting that adaptive increases in thermogenic capacity involve integrated functional changes across the O_2 cascade. VO_{2max} in hypoxia appears to be enhanced in high-altitude deer mice (Fig. 1) *via* increases in pulmonary O_2 uptake (Fig. 2A), haemoglobin- O_2 affinity (Snyder et al., 1982; Chappell and Snyder, 1984; Storz et al., 2010a; Natarajan et al., 2013), cardiac performance and/or blood O_2 extraction (Fig. 2C), and the capacity for O_2 diffusion and utilization in skeletal muscle (Cheviron et al., 2012; Cheviron et al., 2014; Lui et al., 2015; Scott et al., 2015; Mahalingam et al., 2017). Therefore, the concerted evolution of physiological systems underlying O_2 transport and utilization appear critical to the process of high-altitude adaptation.

ACKNOWLEDGEMENTS

The authors would like to thank Paras Patel and Oliver Wearing for technical assistance with data collection, and two anonymous referees for helpful comments on an earlier version of this manuscript.

COMPETING INTERESTS

The authors declare no competing or financial interests.

AUTHOR CONTRIBUTIONS

G.R.S. designed the experiments. K.B.T. led and all other authors contributed to data collection and/or analysis. K.B.T. and G.R.S. drafted the manuscript. All authors contributed to interpreting the data and revising the manuscript, and all gave final approval for publication.

FUNDING

This research was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant to G.R.S., National Science Foundation grants to Z.A.C. (IOS-1354934 and IOS-1634219) and J.F.S. (IOS-1354390), and a National Institutes of Health grant to J.F.S. (HL087216). C.M.I. was supported by a NSERC Postgraduate Scholarship and an Ontario Graduate Scholarship. G.R.S. is supported by the Canada Research Chairs Program.

REFERENCES

- Brutsaert, T.** (2016). Why are high altitude natives so strong at high altitude? Nature vs. nurture: genetic factors vs. growth and development. *Adv. Exp. Med. Biol.* **903**, 101-112.
- Chappell, M. A. and Hammond, K. A.** (2004). Maximal aerobic performance of deer mice in combined cold and exercise challenges. *J. Comp. Physiol. B* **174**, 41-48.
- Chappell, M. A., Hammond, K. A., Cardullo, R. A., Russell, G. A., Rezende, E. L. and Miller, C.** (2007). Deer mouse aerobic performance across altitudes: effects of developmental history and temperature acclimation. *Physiol. Biochem. Zool.* **80**, 652-662.
- Chappell, M. A., Hayes, J. P. and Lee, R. G. S.** (1988). Hemoglobin polymorphisms in deer mice (*Peromyscus maniculatus*): physiology of beta-globin variants and alpha-globin recombinants. *Evolution* **42**, 681-688.
- Chappell, M. A. and Snyder, L. R. G.** (1984). Biochemical and physiological correlates of deer mouse α -chain hemoglobin polymorphisms. *Proc. Natl. Acad. Sci. U. S. A.* **81**, 5484-5488.
- Chevron, Z. A., Bachman, G. C., Connaty, A. D., McClelland, G. B. and Storz, J. F.** (2012). Regulatory changes contribute to the adaptive enhancement of thermogenic capacity in high-altitude deer mice. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8635-8640.
- Chevron, Z. A., Bachman, G. C. and Storz, J. F.** (2013). Contributions of phenotypic plasticity to differences in thermogenic performance between highland and lowland deer mice. *J. Exp. Biol.* **216**, 1160-1166.
- Chevron, Z. A., Connaty, A. D., McClelland, G. B. and Storz, J. F.** (2014). Functional genomics of adaptation to hypoxic cold-stress in high-altitude deer mice: transcriptomic plasticity and thermogenic performance. *Evolution* **68**, 48-62.
- Dalziel, A. C., Rogers, S. M. and Schulte, P. M.** (2009). Linking genotypes to phenotypes and fitness: how mechanistic biology can inform molecular ecology. *Mol. Ecol.* **18**, 4997-5017.
- Dawson, N. J., Ivy, C. M., Alza, L., Cheek, R., York, J. M., Chua, B., Milsom, W. K., McCracken, K. G. and Scott, G. R.** (2016). Mitochondrial physiology in the skeletal and cardiac muscles is altered in torrent ducks, *Merganetta armata*, from high altitudes in the Andes. *J. Exp. Biol.* **219**, 3719-3728.

- Garland, T. J. and Carter, P. A.** (1994). Evolutionary physiology. *Annu. Rev. Physiol.* **56**, 579-621.
- Hammond, K. A., Chappell, M. A. and Kristan, D. M.** (2002). Developmental plasticity in aerobic performance in deer mice (*Peromyscus maniculatus*). *Comparative Biochemistry and Physiology a-Molecular and Integrative Physiology* **133**, 213-224.
- Hammond, K. A., Szewczak, J. and Krol, E.** (2001). Effects of altitude and temperature on organ phenotypic plasticity along an altitudinal gradient. *J. Exp. Biol.* **204**, 1991-2000.
- Hayes, J. P.** (1989a). Altitudinal and seasonal effects on aerobic metabolism of deer mice. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology* **159**, 453-459.
- Hayes, J. P.** (1989b). Field and maximal metabolic rates of deer mice (*Peromyscus maniculatus*) at low and high altitudes. *Physiol. Zool.* **62**, 732-744.
- Hayes, J. P. and O'Connor, C. S.** (1999). Natural selection on thermogenic capacity of high-altitude deer mice. *Evolution* **53**, 1280-1287.
- Hock, R. J.** (1964). Physiological responses of deer mice to various native altitudes. In *The Physiological Effects of High Altitude*, (ed. W. H. Weihe), pp. 59-72. New York: Macmillan.
- Ivy, C. M. and Scott, G. R.** (2015). Control of breathing and the circulation in high-altitude mammals and birds. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **186**, 66-74.
- Ivy, C. M. and Scott, G. R.** (2017). Control of breathing and ventilatory acclimatization to hypoxia in deer mice native to high altitudes. *Acta Physiol.* **In press**.
- Lau, D. S., Connaty, A. D., Mahalingam, S., Wall, N., Cheviron, Z. A., Storz, J. F., Scott, G. R. and McClelland, G. B.** (2017). Acclimation to hypoxia increases carbohydrate use during exercise in high-altitude deer mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **312**, R400-R411.
- Lighton, J. R. B.** (2008). Measuring metabolic rates: a manual for scientists: Oxford University Press, USA.
- Lui, M. A., Mahalingam, S., Patel, P., Connaty, A. D., Ivy, C. M., Cheviron, Z. A., Storz, J. F., McClelland, G. B. and Scott, G. R.** (2015). High-altitude ancestry and hypoxia acclimation have distinct effects on exercise capacity and muscle phenotype in deer mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **308**, R779-R791.

- Mahalingam, S., McClelland, G. B. and Scott, G. R.** (2017). Evolved changes in the intracellular distribution and physiology of muscle mitochondria in high-altitude native deer mice. *J. Physiol. (Lond)*. **Accepted**, doi:10.1113/JP274130.
- Monge, C. and León-Velarde, F.** (1991). Physiological adaptation to high altitude: oxygen transport in mammals and birds. *Physiol. Rev.* **71**, 1135-1172.
- Natarajan, C., Hoffmann, F. G., Lanier, H. C., Wolf, C. J., Cheviron, Z. A., Spangler, M. L., Weber, R. E., Fago, A. and Storz, J. F.** (2015). Intraspecific polymorphism, interspecific divergence, and the origins of function-altering mutations in deer mouse hemoglobin. *Mol. Biol. Evol.* **32**, 978-997.
- Natarajan, C., Inoguchi, N., Weber, R. E., Fago, A., Moriyama, H. and Storz, J. F.** (2013). Epistasis among adaptive mutations in deer mouse hemoglobin. *Science* **340**, 1324-1327.
- R Core Team.** (2016). R: A language and environment for statistical computing. In *R Foundation for Statistical Computing* Vienna, Austria: (<http://www.R-project.org/>).
- Rosenmann, M. and Morrison, P.** (1974). Maximum oxygen consumption and heat loss facilitation in small homeotherms by He-O₂. *Am. J. Physiol.* **226**, 490-495.
- Russell, G. A., Rezende, E. L. and Hammond, K. A.** (2008). Development partly determines the aerobic performance of adult deer mice, *Peromyscus maniculatus*. *J. Exp. Biol.* **211**, 35-41.
- Scott, G. R.** (2011). Elevated performance: the unique physiology of birds that fly at high altitudes. *J. Exp. Biol.* **214**, 2455-2462.
- Scott, G. R., Elogio, T. S., Lui, M. A., Storz, J. F. and Cheviron, Z. A.** (2015). Adaptive modifications of muscle phenotype in high-altitude deer mice are associated with evolved changes in gene regulation. *Mol. Biol. Evol.* **32**, 1962-1976.
- Scott, G. R., Schulte, P. M., Egginton, S., Scott, A. L. M., Richards, J. G. and Milsom, W. K.** (2011). Molecular evolution of cytochrome c oxidase underlies high-altitude adaptation in the bar-headed goose. *Mol. Biol. Evol.* **28**, 351-363.
- Sheafor, B. A.** (2003). Metabolic enzyme activities across an altitudinal gradient: an examination of pikas (genus *Ochotona*). *J. Exp. Biol.* **206**, 1241-1249.
- Shirkey, N. J. and Hammond, K. A.** (2014). The relationship between cardiopulmonary size and aerobic performance in adult deer mice at high altitude. *J. Exp. Biol.* **217**, 3758-3764.

- Snyder, L. R. G., Born, S. and Lechner, A. J.** (1982). Blood oxygen affinity in high- and low-altitude populations of the deer mouse. *Respir. Physiol.* **48**, 89-105.
- Storz, J. F., Runck, A. M., Moriyama, H., Weber, R. E. and Fago, A.** (2010a). Genetic differences in hemoglobin function between highland and lowland deer mice. *J. Exp. Biol.* **213**, 2565-2574.
- Storz, J. F., Scott, G. R. and Cheviron, Z. A.** (2010b). Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. *J. Exp. Biol.* **213**, 4125-4136.
- Velotta, J. P., Jones, J., Wolf, C. J. and Cheviron, Z. A.** (2016). Transcriptomic plasticity in brown adipose tissue contributes to an enhanced capacity for nonshivering thermogenesis in deer mice. *Mol. Ecol.* **25**, 2870-2886.

Figures




Fig. 1. Thermogenic capacity, measured in hypoxia as the maximal rate of O_2 consumption (VO_{2max}) during acute cold exposure, was enhanced in high-altitude deer mice. The effects of population altitude and hypoxia acclimation on hypoxic VO_{2max} were assessed by calculating the residuals (A) for an allometric regression of hypoxic VO_{2max} to body mass (M_b ; $VO_{2max}=0.46 M_b^{0.65}$) (B). (A) The left axis shows the hypoxic VO_{2max} for an average-sized 21.6-g mouse, calculated for each group by adding the residual (shown on the right axis) to the VO_{2max} predicted at 21.6 g by the regression (means \pm s.e.m.). There were significant main effects of population (*, $F_{1,36}=4.42$, $p=0.043$) and hypoxia acclimation (\dagger , $F_{1,36}=18.25$, $p<0.001$). (B) Dashed grey lines represent 95% confidence intervals of the allometric regression (∇ , normoxic highlanders, $n=15$; ∇ , hypoxic highlanders, $n=14$; \blacktriangle , normoxic lowlanders, $n=6$; \triangle , hypoxic lowlanders, $n=5$).




Fig. 2. Population altitude and hypoxia acclimation affect circulatory O₂ delivery at hypoxic VO₂max in deer mice. (A) Arterial O₂ saturation was higher in highland mice (*, $F_{1,24}=10.46$, $p=0.004$) but was unaffected by hypoxia acclimation ($F_{1,24}=0.063$, $p=0.803$). (B) Heart rate at VO₂max increased after hypoxia acclimation (†, $F_{1,30}=15.48$, $p<0.001$) but was similar between populations ($F_{1,30}=0.129$, $p=0.722$). The amount of O₂ extracted from the blood per heartbeat ('O₂ pulse', quotient of VO₂ and heart rate), assessed by calculating the residuals (C) for an allometric regression to body mass (M_b ; O₂ pulse = $1.15 M_b^{0.51}$) (D) was greater in highland mice (*, $F_{1,30}=4.41$, $p=0.044$) and increased after hypoxia acclimation (†, $F_{1,30}=6.10$, $p=0.019$) (see Fig. 1 and Materials and Methods for additional details on this approach). Dashed grey lines in (D) represent 95% confidence intervals of the allometric regression (▼, normoxic highlanders, $n=14$; ▽, hypoxic highlanders, $n=12$; ▲, normoxic lowlanders, $n=4$; △, hypoxic lowlanders, $n=4$). Data are means \pm s.e.m., except in (D) where data from individuals are shown.




Fig. 3. Capillarity of the heart tissue increased after hypoxia acclimation. (A,B)

Representative images of left ventricle tissue near the epicardium, stained for alkaline phosphatase activity to identify capillaries, in normoxia acclimated deer mice from the low-altitude (A) and high-altitude (B) populations (scale bar represents 50 μm). (C) There was a significant main effect of hypoxia acclimation on capillary density (†, $F_{1,30}=5.53$, $p=0.026$), but no difference between populations ($F_{1,30}=0.57$, $p=0.455$). Data are means \pm s.e.m., with sample sizes as follows: normoxia acclimated highlanders, N=9; hypoxia acclimated highlanders, N=8; normoxia acclimated lowlanders, N=7; hypoxia acclimated lowlanders, N=10.

Table S1. Statistical results from linear models used to examine the effects of body mass, population altitude, and acclimation environment on VO₂max and O₂ pulse

Source	df	F	Probability > F
VO₂max			
Mass	1	21.6	<0.001
Population altitude	1	4.22	0.048
Acclimation environment	1	27.8	<0.001
Altitude × acclimation	1	0.218	0.644
Residual	35		
O₂ pulse			
Mass	1	13.0	0.001
Population altitude	1	3.87	0.059
Acclimation environment	1	12.2	0.002
Altitude × acclimation	1	0.374	0.546
Residual	29		

df, degrees of freedom; O₂ pulse, the amount of O₂ extracted from the blood per heartbeat (the quotient of VO₂ and f_H); VO₂max, maximal O₂ consumption measured during cold exposure in a hypoxic heliox atmosphere (12% O₂, 88% He).

Body mass (Mb) and the variable of interest were log-transformed before making statistical comparisons, which were carried out using linear models (lm) in R (LogVariable ~ LogMb + Altitude + Acclimation + Altitude×Acclimation).